

Climate indices with CDO

Climate indices of daily temperature and precipitation extremes
October 2015

Uwe Schulzweida – *MPI for Meteorology*

Ralf Quast – *Brockmann Consult*

Contents

1	Introduction	3
2	Climate indices reference manual	4
2.0.1	ECACDD - Consecutive dry days index per time period	6
2.0.2	ECACFD - Consecutive frost days index per time period	6
2.0.3	ECACSU - Consecutive summer days index per time period	7
2.0.4	ECACWD - Consecutive wet days index per time period	7
2.0.5	ECACWDI - Cold wave duration index w.r.t. mean of reference period	8
2.0.6	ECAGWFI - Cold-spell days index w.r.t. 10th percentile of reference period	8
2.0.7	ECAETR - Intra-period extreme temperature range	10
2.0.8	ECAFD - Frost days index per time period	10
2.0.9	ECAFSL - Thermal Growing season length index	11
2.0.10	ECAHD - Heating degree days per time period	12
2.0.11	ECAHWDI - Heat wave duration index w.r.t. mean of reference period	12
2.0.12	ECAHWFI - Warm spell days index w.r.t. 90th percentile of reference period	13
2.0.13	ECAID - Ice days index per time period	13
2.0.14	ECAR75P - Moderate wet days w.r.t. 75th percentile of reference period	14
2.0.15	ECAR75PTOT - Precipitation percent due to R75p days	14
2.0.16	ECAR90P - Wet days w.r.t. 90th percentile of reference period	15
2.0.17	ECAR90PTOT - Precipitation percent due to R90p days	15
2.0.18	ECAR95P - Very wet days w.r.t. 95th percentile of reference period	16
2.0.19	ECAR95PTOT - Precipitation percent due to R95p days	16
2.0.20	ECAR99P - Extremely wet days w.r.t. 99th percentile of reference period	17
2.0.21	ECAR99PTOT - Precipitation percent due to R99p days	17
2.0.22	ECAPD - Precipitation days index per time period	18
2.0.23	ECARR1 - Wet days index per time period	19
2.0.24	ECARX1DAY - Highest one day precipitation amount per time period	19
2.0.25	ECARX5DAY - Highest five-day precipitation amount per time period	21
2.0.26	ECASDII - Simple daily intensity index per time period	21
2.0.27	ECASU - Summer days index per time period	22
2.0.28	ECATG10P - Cold days percent w.r.t. 10th percentile of reference period	23
2.0.29	ECATG90P - Warm days percent w.r.t. 90th percentile of reference period	23
2.0.30	ECATN10P - Cold nights percent w.r.t. 10th percentile of reference period	24
2.0.31	ECATN90P - Warm nights percent w.r.t. 90th percentile of reference period	24
2.0.32	ECATR - Tropical nights index per time period	25
2.0.33	ECATX10P - Very cold days percent w.r.t. 10th percentile of reference period	25
2.0.34	ECATX90P - Very warm days percent w.r.t. 90th percentile of reference period	26
	Operator index	28

1 Introduction

The Climate Data Operators (**CDO**) software is a collection of operators for standard processing of climate and forecast model data.

This document describes additional **CDO** operators to compute climate indices of daily temperature and precipitation extreme. The definition of these climate indices are from the European Climate Assessment (ECA) project.

The climate indices were implemented in **CDO** by Ralf Quast (Brockmann Consult) on behalf of the Service Gruppe Anpassung (SGA) in 2006. SGA was part of the Model and Data Group (M&D) at the MPI for Meteorology. In 2010, the Model and Data Group became the Data Management department at DKRZ (Deutsches Klimarechenzentrum) and the SGA was disintegrated. For this reason there is no further user support available for these **CDO** operators.

2 Climate indices reference manual

This section gives a description of all **CDO** operators to compute the climate indices of daily temperature and precipitation extreme. Related operators are grouped to modules. For easier description all single input files are named `infile` or `infile1`, `infile2`, etc., and an arbitrary number of input files are named `ifiles`. All output files are named `ofile` or `ofile1`, `ofile2`, etc. Further the following notion is introduced:

- $i(t)$ Timestep t of `infile`
- $i(t, x)$ Element number x of the field at timestep t of `infile`
- $o(t)$ Timestep t of `ofile`
- $o(t, x)$ Element number x of the field at timestep t of `ofile`

Here is a short overview of all operators in this section:

eca_cdd	Consecutive dry days index per time period
eca_cfd	Consecutive frost days index per time period
eca_csu	Consecutive summer days index per time period
eca_cwd	Consecutive wet days index per time period
eca_cwdi	Cold wave duration index w.r.t. mean of reference period
eca_cwfi	Cold-spell days index w.r.t. 10th percentile of reference period
eca_etr	Intra-period extreme temperature range
eca_fd	Frost days index per time period
eca_gsl	Growing season length index
eca_hd	Heating degree days per time period
eca_hwdi	Heat wave duration index w.r.t. mean of reference period
eca_hwfi	Warm spell days index w.r.t. 90th percentile of reference period
eca_id	Ice days index per time period
eca_r75p	Moderate wet days w.r.t. 75th percentile of reference period
eca_r75ptot	Precipitation percent due to R75p days
eca_r90p	Wet days w.r.t. 90th percentile of reference period
eca_r90ptot	Precipitation percent due to R90p days
eca_r95p	Very wet days w.r.t. 95th percentile of reference period
eca_r95ptot	Precipitation percent due to R95p days

eca_r99p	Extremely wet days w.r.t. 99th percentile of reference period
eca_r99ptot	Precipitation percent due to R99p days
eca_pd	Precipitation days index per time period
eca_r10mm	Heavy precipitation days index per time period
eca_r20mm	Very heavy precipitation days index per time period
eca_rr1	Wet days index per time period
eca_rx1day	Highest one day precipitation amount per time period
eca_rx5day	Highest five-day precipitation amount per time period
eca_sdii	Simple daily intensity index per time period
eca_su	Summer days index per time period
eca_tg10p	Cold days percent w.r.t. 10th percentile of reference period
eca_tg90p	Warm days percent w.r.t. 90th percentile of reference period
eca_tn10p	Cold nights percent w.r.t. 10th percentile of reference period
eca_tn90p	Warm nights percent w.r.t. 90th percentile of reference period
eca_tr	Tropical nights index per time period
eca_tx10p	Very cold days percent w.r.t. 10th percentile of reference period
eca_tx90p	Very warm days percent w.r.t. 90th percentile of reference period

2.0.1 ECACDD - Consecutive dry days index per time period

Synopsis

```
eca_cdd[,R] ifile ofile
```

Description

Let `ifile` be a time series of the daily precipitation amount `RR`, then the largest number of consecutive days where `RR` is less than `R` is counted. `R` is an optional parameter with default `R = 1` mm. A further output variable is the number of dry periods of more than 5 days. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- consecutive_dry_days_index_per_time_period
- number_of_cdd_periods_with_more_than_5days_per_time_period

Parameter

`R` FLOAT Precipitation threshold (unit: mm; default: `R = 1` mm)

Example

To get the largest number of consecutive dry days of a time series of daily precipitation amounts use:

```
cdo eca_cdd rrfile ofile
```

2.0.2 ECACFD - Consecutive frost days index per time period

Synopsis

```
eca_cfd ifile ofile
```

Description

Let `ifile` be a time series of the daily minimum temperature `TN`, then the largest number of consecutive days where `TN` < 0 °C is counted. Note that `TN` have to be given in units of Kelvin. A further output variable is the number of frost periods of more than 5 days. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- consecutive_frost_days_index_per_time_period
- number_of_cfd_periods_with_more_than_5days_per_time_period

Example

To get the largest number of consecutive frost days of a time series of daily minimum temperatures use:

```
cdo eca_cfd tnfile ofile
```

2.0.3 ECACSU - Consecutive summer days index per time period

Synopsis

```
eca_csu[,T] ifile ofile
```

Description

Let `ifile` be a time series of the daily maximum temperature TX, then the largest number of consecutive days where $TX > T$ is counted. The number T is an optional parameter with default $T = 25^{\circ}\text{C}$. Note that TN have to be given in units of Kelvin, whereas T have to be given in degrees Celsius. A further output variable is the number of summer periods of more than 5 days. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- consecutive_summer_days_index_per_time_period
- number_of_csu_periods_with_more_than_5days_per_time_period

Parameter

`T` FLOAT Temperature threshold (unit: $^{\circ}\text{C}$; default: $T = 25^{\circ}\text{C}$)

Example

To get the largest number of consecutive summer days of a time series of daily maximum temperatures use:

```
cdo eca_csu txfile ofile
```

2.0.4 ECACWD - Consecutive wet days index per time period

Synopsis

```
eca_cwd[,R] ifile ofile
```

Description

Let `ifile` be a time series of the daily precipitation amount RR, then the largest number of consecutive days where RR is at least R is counted. R is an optional parameter with default $R = 1$ mm. A further output variable is the number of wet periods of more than 5 days. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- consecutive_wet_days_index_per_time_period
- number_of_cwd_periods_with_more_than_5days_per_time_period

Parameter

`R` FLOAT Precipitation threshold (unit: mm; default: $R = 1$ mm)

Example

To get the largest number of consecutive wet days of a time series of daily precipitation amounts use:

```
cdo eca_cwd rrfile ofile
```

2.0.5 ECACWDI - Cold wave duration index w.r.t. mean of reference period

Synopsis

```
eca_cwdi[,nday[,T]] ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily minimum temperature `TN`, and let `ifile2` be the mean `TNnorm` of daily minimum temperatures for any period used as reference. Then counted is the number of days where, in intervals of at least `nday` consecutive days, $TN < TNnorm - T$. The numbers `nday` and `T` are optional parameters with default `nday = 6` and `T = 5°C`. A further output variable is the number of cold waves longer than or equal to `nday` days. `TNnorm` is calculated as the mean of minimum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both `TN` and `TNnorm` have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- cold_wave_duration_index_wrt_mean_of_reference_period
- cold_waves_per_time_period

Parameter

<code>nday</code>	INTEGER	Number of consecutive days (default: <code>nday = 6</code>)
<code>T</code>	FLOAT	Temperature offset (unit: °C; default: <code>T = 5°C</code>)

Example

To compute the cold wave duration index of a time series of daily minimum temperatures use:

```
cdo eca_cwdi tnfile tnnormfile ofile
```

2.0.6 ECACWFI - Cold-spell days index w.r.t. 10th percentile of reference period

Synopsis

```
eca_cwfi[,nday] ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily mean temperature `TG`, and `ifile2` be the 10th percentile `TGn10` of daily mean temperatures for any period used as reference. Then counted is the number of days where, in intervals of at least `nday` consecutive days, $TG < TGn10$. The number `nday` is an optional parameter with default `nday = 6`. A further output variable is the number of cold-spell periods longer than or equal to `nday` days. `TGn10` is calculated as the 10th percentile of daily mean temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both `TG` and `TGn10` have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- cold_spell_days_index_wrt_10th_percentile_of_reference_period
- cold_spell_periods_per_time_period

Parameter

nday INTEGER Number of consecutive days (default: *nday* = 6)

Example

To compute the number of cold-spell days of a time series of daily mean temperatures use:

```
cdo eca_cwfi tgfile tgn10file ofile
```

2.0.7 ECAETR - Intra-period extreme temperature range

Synopsis

```
eca_etr ifile1 ifile2 ofile
```

Description

Let `ifile1` and `ifile2` be time series of the maximum and minimum temperature TX and TN, respectively. Then the extreme temperature range is the difference of the maximum of TX and the minimum of TN. Note that TX and TN have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timesteps in `ifile1` and `ifile2`. The following variables are created:

- `intra_period_extreme_temperature_range`

Example

To get the intra-period extreme temperature range for two time series of maximum and minimum temperatures use:

```
cdo eca_etr txfile tnfile ofile
```

2.0.8 ECAFD - Frost days index per time period

Synopsis

```
eca_fd ifile ofile
```

Description

Let `ifile` be a time series of the daily minimum temperature TN, then the number of days where $TN < 0\text{ }^{\circ}\text{C}$ is counted. Note that TN have to be given in units of Kelvin. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `frost_days_index_per_time_period`

Example

To get the number of frost days of a time series of daily minimum temperatures use:

```
cdo eca_fd tnfile ofile
```

2.0.9 ECAGSL - Thermal Growing season length index

Synopsis

```
eca_gsl[,nday[,T[,fland]]] ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily mean temperature TG, and `ifile2` be a land-water mask. Within a period of 12 months, the thermal growing season length is officially defined as the number of days between:

- first occurrence of at least `nday` consecutive days with $TG > T$
- first occurrence of at least `nday` consecutive days with $TG < T$ within the last 6 months

On northern hemispere, this period corresponds with the regular year, whereas on southern hemispere, it starts at July 1st. Please note, that this definition may lead to weird results concerning values $TG = T$: In the first half of the period, these days do not contribute to the gsl, but they do within the second half. Moreover this definition could lead to discontinuous values in equatorial regions.

The numbers `nday` and `T` are optional parameter with default `nday = 6` and `T = 5°C`. The number `fland` is an optional parameter with default value `fland = 0.5` and denotes the fraction of a grid point that have to be covered by land in order to be included in the calculation. A further output variable is the start day of year of the growing season. Note that TG have to be given in units of Kelvin, whereas `T` have to be given in degrees Celsius.

The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `thermal_growing_season_length`
- `day_of_year_of_growing_season_start`

Parameter

<code>nday</code>	INTEGER	Number of consecutive days (default: <code>nday = 6</code>)
<code>T</code>	FLOAT	Temperature threshold (unit: °C; default: <code>T = 5°C</code>)
<code>fland</code>	FLOAT	Land fraction threshold (default: <code>fland = 0.5</code>)

Example

To get the growing season length of a time series of daily mean temperatures use:

```
cdo eca_gsl tgfile maskfile ofile
```

2.0.10 ECAHD - Heating degree days per time period

Synopsis

```
eca_hd[,T1[,T2]] ifile ofile
```

Description

Let `ifile` be a time series of the daily mean temperature `TG`, then the heating degree days are defined as the sum of $T1 - TG$, where only values $TG < T2$ are considered. If $T1$ and $T2$ are omitted, a temperature of 17°C is used for both parameters. If only $T1$ is given, $T2$ is set to $T1$. Note that `TG` have to be given in units of kelvin, whereas $T1$ and $T2$ have to be given in degrees Celsius. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `heating_degree_days_per_time_period`

Parameter

$T1$	FLOAT	Temperature limit (unit: °C; default: $T1 = 17^{\circ}\text{C}$)
$T2$	FLOAT	Temperature limit (unit: °C; default: $T2 = T1$)

Example

To compute the heating degree days of a time series of daily mean temperatures use:

```
cdo eca_hd tgfile ofile
```

2.0.11 ECAHWDI - Heat wave duration index w.r.t. mean of reference period

Synopsis

```
eca_hwdi[,nday[,T]] ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily maximum temperature `TX`, and let `ifile2` be the mean `TXnorm` of daily maximum temperatures for any period used as reference. Then counted is the number of days where, in intervals of at least `nday` consecutive days, $TX > TXnorm + T$. The numbers `nday` and T are optional parameters with default `nday = 6` and $T = 5^{\circ}\text{C}$. A further output variable is the number of heat waves longer than or equal to `nday` days. `TXnorm` is calculated as the mean of maximum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both `TX` and `TXnorm` have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `heat_wave_duration_index_wrt_mean_of_reference_period`
- `heat_waves_per_time_period`

Parameter

<code>nday</code>	INTEGER	Number of consecutive days (default: <code>nday = 6</code>)
T	FLOAT	Temperature offset (unit: °C; default: $T = 5^{\circ}\text{C}$)

2.0.12 ECAHWFI - Warm spell days index w.r.t. 90th percentile of reference period

Synopsis

```
eca_hwfi[,nday] ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily mean temperature TG, and `ifile2` be the 90th percentile TGn90 of daily mean temperatures for any period used as reference. Then counted is the number of days where, in intervals of at least *nday* consecutive days, $TG > TGn90$. The number *nday* is an optional parameter with default *nday* = 6. A further output variable is the number of warm-spell periods longer than or equal to *nday* days. TGn90 is calculated as the 90th percentile of daily mean temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TG and TGn90 have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- warm_spell_days_index_wrt_90th_percentile_of_reference_period
- warm_spell_periods_per_time_period

Parameter

nday INTEGER Number of consecutive days (default: *nday* = 6)

Example

To compute the number of warm-spell days of a time series of daily mean temperatures use:

```
cdo eca_hwfi tgfile tgn90file ofile
```

2.0.13 ECAID - Ice days index per time period

Synopsis

```
eca_id ifile ofile
```

Description

Let `ifile` be a time series of the daily maximum temperature TX, then the number of days where $TX < 0$ °C is counted. Note that TX have to be given in units of Kelvin. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- ice_days_index_per_time_period

Example

To get the number of ice days of a time series of daily maximum temperatures use:

```
cdo eca_id txfile ofile
```

2.0.14 ECAR75P - Moderate wet days w.r.t. 75th percentile of reference period

Synopsis

```
eca_r75p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 75th percentile RRn75 of the daily precipitation amount at wet days for any period used as reference. Then the percentage of wet days with $RR > RRn75$ is calculated. RRn75 is calculated as the 75th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypct1,75`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- moderate_wet_days_wrt_75th_percentile_of_reference_period

Example

To compute the percentage of wet days with daily precipitation amount greater than the 75th percentile of the daily precipitation amount at wet days for a given reference period use:

```
cdo eca_r75p rrfile rrn75file ofile
```

2.0.15 ECAR75PTOT - Precipitation percent due to R75p days

Synopsis

```
eca_r75ptot ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 75th percentile RRn75 of the daily precipitation amount at wet days for any period used as reference. Then the ratio of the precipitation sum at wet days with $RR > RRn75$ to the total precipitation sum is calculated. RRn75 is calculated as the 75th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypct1,75`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- precipitation_percent_due_to_R75p_days

2.0.16 ECAR90P - Wet days w.r.t. 90th percentile of reference period

Synopsis

```
eca_r90p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 90th percentile RRn90 of the daily precipitation amount at wet days for any period used as reference. Then the percentage of wet days with $RR > RRn90$ is calculated. RRn90 is calculated as the 90th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypctl,90`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `wet_days_wrt_90th_percentile_of_reference_period`

Example

To compute the percentage of wet days where the daily precipitation amount is greater than the 90th percentile of the daily precipitation amount at wet days for a given reference period use:

```
cdo eca_r90p rrfile rrn90file ofile
```

2.0.17 ECAR90PTOT - Precipitation percent due to R90p days

Synopsis

```
eca_r90ptot ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 90th percentile RRn90 of the daily precipitation amount at wet days for any period used as reference. Then the ratio of the precipitation sum at wet days with $RR > RRn90$ to the total precipitation sum is calculated. RRn90 is calculated as the 90th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypctl,90`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `precipitation_percent_due_to_R90p_days`

2.0.18 ECAR95P - Very wet days w.r.t. 95th percentile of reference period

Synopsis

```
eca_r95p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 95th percentile RRn95 of the daily precipitation amount at wet days for any period used as reference. Then the percentage of wet days with $RR > RRn95$ is calculated. RRn95 is calculated as the 95th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypct1,95`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `very_wet_days_wrt_95th_percentile_of_reference_period`

Example

To compute the percentage of wet days where the daily precipitation amount is greater than the 95th percentile of the daily precipitation amount at wet days for a given reference period use:

```
cdo eca_r95p rrfile rrn95file ofile
```

2.0.19 ECAR95PTOT - Precipitation percent due to R95p days

Synopsis

```
eca_r95ptot ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 95th percentile RRn95 of the daily precipitation amount at wet days for any period used as reference. Then the ratio of the precipitation sum at wet days with $RR > RRn95$ to the total precipitation sum is calculated. RRn95 is calculated as the 95th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypct1,95`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `precipitation_percent_due_to_R95p_days`

2.0.20 ECAR99P - Extremely wet days w.r.t. 99th percentile of reference period

Synopsis

```
eca_r99p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 99th percentile RRn99 of the daily precipitation amount at wet days for any period used as reference. Then the percentage of wet days with $RR > RRn99$ is calculated. RRn99 is calculated as the 99th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypctl,99`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `extremely_wet_days_wrt_99th_percentile_of_reference_period`

Example

To compute the percentage of wet days where the daily precipitation amount is greater than the 99th percentile of the daily precipitation amount at wet days for a given reference period use:

```
cdo eca_r99p rrfile rrn99file ofile
```

2.0.21 ECAR99PTOT - Precipitation percent due to R99p days

Synopsis

```
eca_r99ptot ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series RR of the daily precipitation amount at wet days (precipitation ≥ 1 mm) and `ifile2` be the 99th percentile RRn99 of the daily precipitation amount at wet days for any period used as reference. Then the ratio of the precipitation sum at wet days with $RR > RRn99$ to the total precipitation sum is calculated. RRn99 is calculated as the 99th percentile of all wet days of a given climate reference period. Usually `ifile2` is generated by the operator `ydaypctl,99`. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `precipitation_percent_due_to_R99p_days`

2.0.22 ECAPD - Precipitation days index per time period

Synopsis

```
eca_pd,x ifile ofile
eca_r10mm ifile ofile
eca_r20mm ifile ofile
```

Description

Let `ifile` be a time series of the daily precipitation amount `RR` in [mm] (or alternatively in [kg m⁻²]), then the number of days where `RR` is at least x mm is counted. `eca_r10mm` and `eca_r20mm` are specific ECA operators with a daily precipitation amount of 10 and 20 mm respectively. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `precipitation_days_index_per_time_period`

Operators

<code>eca_pd</code>	Precipitation days index per time period Generic ECA operator with daily precipitation sum exceeding x mm.
<code>eca_r10mm</code>	Heavy precipitation days index per time period Specific ECA operator with daily precipitation sum exceeding 10 mm.
<code>eca_r20mm</code>	Very heavy precipitation days index per time period Specific ECA operator with daily precipitation sum exceeding 20 mm.

Parameter

<code>x</code>	FLOAT	Daily precipitation amount threshold in [mm]
----------------	-------	--

Note

Precipitation rates in [mm/s] have to be converted to precipitation amounts (multiply with 86400 s). Apart from metadata information the result of `eca_pd,1` and `eca_rr1` is the same.

Example

To get the number of days with precipitation greater than 25 mm for a time series of daily precipitation amounts use:

```
cdo eca_pd,25 ifile ofile
```

2.0.23 ECARR1 - Wet days index per time period

Synopsis

```
eca_rr1[,R] ifile ofile
```

Description

Let *ifile* be a time series of the daily precipitation amount *RR* in [mm] (or alternatively in [kg m⁻²]), then the number of days where *RR* is at least *R* is counted. *R* is an optional parameter with default *R* = 1 mm. The date information of a timestep in *ofile* is the date of the last contributing timestep in *ifile*. The following variables are created:

- wet_days_index_per_time_period

Parameter

R FLOAT Precipitation threshold (unit: mm; default: *R* = 1 mm)

Example

To get the number of wet days of a time series of daily precipitation amounts use:

```
cdo eca_rr1 rrfile ofile
```

2.0.24 ECARX1DAY - Highest one day precipitation amount per time period

Synopsis

```
eca_rx1day[,mode] ifile ofile
```

Description

Let *ifile* be a time series of the daily precipitation amount *RR*, then the maximum of *RR* is written to *ofile*. If the optional parameter *mode* is set to 'm' the maximum daily precipitation amounts are determined for each month. The date information of a timestep in *ofile* is the date of the last contributing timestep in *ifile*. The following variables are created:

- highest_one_day_precipitation_amount_per_time_period

Parameter

mode STRING Operation mode (optional). If *mode* = 'm' then maximum daily precipitation amounts are determined for each month

Example

To get the maximum of a time series of daily precipitation amounts use:

```
cdo eca_rx1day rrfile ofile
```

If you are interested in the maximum daily precipitation for each month, use:

```
cdo eca_rx1day,m rrfile ofile
```

Apart from metadata information, both operations yield the same as:

```
cdo timmax rrfile ofile  
cdo monmax rrfile ofile
```

2.0.25 ECARX5DAY - Highest five-day precipitation amount per time period

Synopsis

```
eca_rx5day[,x] ifile ofile
```

Description

Let `ifile` be a time series of 5-day precipitation totals `RR`, then the maximum of `RR` is written to `ofile`. A further output variable is the number of 5 day period with precipitation totals greater than x mm, where x is an optional parameter with default $x = 50$ mm. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `highest_five_day_precipitation_amount_per_time_period`
- `number_of_5day_heavy_precipitation_periods_per_time_period`

Parameter

<code>x</code>	FLOAT	Precipitation threshold (unit: mm; default: $x = 50$ mm)
----------------	-------	--

Example

To get the maximum of a time series of 5-day precipitation totals use:

```
cdo eca_rx5day rrfile ofile
```

Apart from metadata information, the above operation yields the same as:

```
cdo timmax rrfile ofile
```

2.0.26 ECASDII - Simple daily intensity index per time period

Synopsis

```
eca_sdii[,R] ifile ofile
```

Description

Let `ifile` be a time series of the daily precipitation amount `RR`, then the mean precipitation amount at wet days ($RR > R$) is written to `ofile`. R is an optional parameter with default $R = 1$ mm. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `simple_daily_intensity_index_per_time_period`

Parameter

<code>R</code>	FLOAT	Precipitation threshold (unit: mm; default: $R = 1$ mm)
----------------	-------	---

Example

To get the daily intensity index of a time series of daily precipitation amounts use:

```
cdo eca_sdii rrfile ofile
```

2.0.27 ECASU - Summer days index per time period

Synopsis

```
eca_su[,T] ifile ofile
```

Description

Let `ifile` be a time series of the daily maximum temperature TX, then the number of days where $TX > T$ is counted. The number T is an optional parameter with default $T = 25^{\circ}\text{C}$. Note that TX have to be given in units of Kelvin, whereas T have to be given in degrees Celsius. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `summer_days_index_per_time_period`

Parameter

T	FLOAT	Temperature threshold (unit: $^{\circ}\text{C}$; default: $T = 25^{\circ}\text{C}$)
-----	-------	---

Example

To get the number of summer days of a time series of daily maximum temperatures use:

```
cdo eca_su txfile ofile
```

2.0.28 ECATG10P - Cold days percent w.r.t. 10th percentile of reference period

Synopsis

```
eca_tg10p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily mean temperature TG, and `ifile2` be the 10th percentile TGn10 of daily mean temperatures for any period used as reference. Then the percentage of time where $TG < TGn10$ is calculated. TGn10 is calculated as the 10th percentile of daily mean temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TG and TGn10 have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `cold_days_percent_wrt_10th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily mean temperature smaller than the 10th percentile of the daily mean temperatures for a given reference period use:

```
cdo eca_tg10p tgfile tgn10file ofile
```

2.0.29 ECATG90P - Warm days percent w.r.t. 90th percentile of reference period

Synopsis

```
eca_tg90p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily mean temperature TG, and `ifile2` be the 90th percentile TGn90 of daily mean temperatures for any period used as reference. Then the percentage of time where $TG > TGn90$ is calculated. TGn90 is calculated as the 90th percentile of daily mean temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TG and TGn90 have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `warm_days_percent_wrt_90th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily mean temperature greater than the 90th percentile of the daily mean temperatures for a given reference period use:

```
cdo eca_tg90p tgfile tgn90file ofile
```

2.0.30 ECATN10P - Cold nights percent w.r.t. 10th percentile of reference period

Synopsis

```
eca_tn10p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily minimum temperature TN, and `ifile2` be the 10th percentile T_{Nn10} of daily minimum temperatures for any period used as reference. Then the percentage of time where $TN < T_{Nn10}$ is calculated. T_{Nn10} is calculated as the 10th percentile of daily minimum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TN and T_{Nn10} have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `cold_nights_percent_wrt_10th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily minimum temperature smaller than the 10th percentile of the daily minimum temperatures for a given reference period use:

```
cdo eca_tn10p tnfile tnn10file ofile
```

2.0.31 ECATN90P - Warm nights percent w.r.t. 90th percentile of reference period

Synopsis

```
eca_tn90p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily minimum temperature TN, and `ifile2` be the 90th percentile T_{Nn90} of daily minimum temperatures for any period used as reference. Then the percentage of time where $TN > T_{Nn90}$ is calculated. T_{Nn90} is calculated as the 90th percentile of daily minimum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TN and T_{Nn90} have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `warm_nights_percent_wrt_90th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily minimum temperature greater than the 90th percentile of the daily minimum temperatures for a given reference period use:

```
cdo eca_tn90p tnfile tnn90file ofile
```


2.0.32 ECATR - Tropical nights index per time period

Synopsis

```
eca_tr[,T] ifile ofile
```

Description

Let `ifile` be a time series of the daily minimum temperature `TN`, then the number of days where `TN` $> T$ is counted. The number T is an optional parameter with default $T = 20^{\circ}\text{C}$. Note that `TN` have to be given in units of Kelvin, whereas T have to be given in degrees Celsius. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile`. The following variables are created:

- `tropical_nights_index_per_time_period`

Parameter

T	FLOAT	Temperature threshold (unit: $^{\circ}\text{C}$; default: $T = 20^{\circ}\text{C}$)
-----	-------	---

Example

To get the number of tropical nights of a time series of daily minimum temperatures use:

```
cdo eca_tr tnfile ofile
```

2.0.33 ECATX10P - Very cold days percent w.r.t. 10th percentile of reference period

Synopsis

```
eca_tx10p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily maximum temperature `TX`, and `ifile2` be the 10th percentile `TXn10` of daily maximum temperatures for any period used as reference. Then the percentage of time where $\text{TX} < \text{TXn10}$ is calculated. `TXn10` is calculated as the 10th percentile of daily maximum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both `TX` and `TXn10` have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `very_cold_days_percent_wrt_10th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily maximum temperature smaller than the 10th percentile of the daily maximum temperatures for a given reference period use:

```
cdo eca_tx10p txfile txn10file ofile
```

2.0.34 ECATX90P - Very warm days percent w.r.t. 90th percentile of reference period

Synopsis

```
eca_tx90p ifile1 ifile2 ofile
```

Description

Let `ifile1` be a time series of the daily maximum temperature TX, and `ifile2` be the 90th percentile TX_{n90} of daily maximum temperatures for any period used as reference. Then the percentage of time where TX > TX_{n90} is calculated. TX_{n90} is calculated as the 90th percentile of daily maximum temperatures of a five day window centred on each calendar day of a given climate reference period. Note that both TX and TX_{n90} have to be given in the same units. The date information of a timestep in `ofile` is the date of the last contributing timestep in `ifile1`. The following variables are created:

- `very_warm_days_percent_wrt_90th_percentile_of_reference_period`

Example

To compute the percentage of timesteps with a daily maximum temperature greater than the 90th percentile of the daily maximum temperatures for a given reference period use:

```
cdo eca_tx90p txfile txn90file ofile
```

Bibliography

[CDI]

[Climate Data Interface](#), from the [Max Planck Institute for Meteorologie](#)

[CDO]

[Climate Data Operators](#), from the [Max Planck Institute for Meteorologie](#)

[ECA]

[ECA indices of extremes](#), from the [Koninklijk Nederlands Meteorologisch Instituut, KNMI](#)

Operator index

E

eca_cdd	6
eca_cfd	6
eca_csu	7
eca_cwd	7
eca_cwdi	8
eca_cwfi	8
eca_etr	10
eca_fd	10
eca_gsl	11
eca_hd	12
eca_hwdi	12
eca_hwfi	13
eca_id	13
eca_pd	18
eca_r10mm	18
eca_r20mm	18
eca_r75p	14
eca_r75ptot	14
eca_r90p	15
eca_r90ptot	15
eca_r95p	16
eca_r95ptot	16
eca_r99p	17
eca_r99ptot	17
eca_rr1	19
eca_rx1day	19
eca_rx5day	21
eca_sdii	21
eca_su	22
eca_tg10p	23
eca_tg90p	23
eca_tn10p	24
eca_tn90p	24
eca_tr	25
eca_tx10p	25
eca_tx90p	26